

**UNIVERSITI TEKNOLOGI MARA**



**INVESTIGATION THE EFFECT OF AP AND PA  
PROJECTIONS ON TESTIS AND OVARY DOSE  
DURING LUMBAR IMAGING AT VARIOUS  
TUBE POTENTIALS (KVP)**

**WAN NUR'IZZATI BT WAN AZIZ**

**BACHELOR IN MEDICAL IMAGING (HONS)  
FACULTY OF HEALTH SCIENCE**

July 2015

## **AUTHOR'S DECLARATION**

I declare that the work in this thesis was carried out in accordance with the regulation of University Teknologi MARA. It is original and is the results of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic institution or non-academic institution for any degree or qualification.

I, hereby, acknowledge that I have been supplied with the Academic Rules and Regulations for Post Graduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

|                      |   |  |
|----------------------|---|--|
| Name of Student      | : | Wan Nur'izzati Bt Wan Aziz   |
| Student I.D. No.     | : | 2011480218   |
| Programme            | : | Bachelor of Medical Imaging (Hons.) – HS222  |
| Faculty              | : | Health and Science   |
| Thesis Title         | : | Investigation the effect of AP and PA projections on testis and ovary dose at various tube potentials (kVp). |
| Signature of Student | : | .....  |
| Date                 | : | July 2015  |

## ABSTRACT

**Purpose:** The aim of this study is to investigate the effect of AP and PA projections on testis and ovary dose at various tube potentials (kVp).

**Methodology:** The study was an experimental study. A whole body and RANDO phantom was exposed three times with different kVp techniques by using computed radiography (CR). The entrance skin doses for testis and ovary were measured by placing the thermoluminescence dosimeters (TLDs). The dose delivered to the testis was measured by create testes using a balloon filled with water and place approximately at location of male testes. Meanwhile, the measurement for ovary dose was done on RANDO phantom at slice 29 and the suitable place for TLD chips placement is 9cm from the anterior surface of abdomen and 6cm from midline based on 'average size' woman. After 24 hour, all TLD chips were read by Harshaw 3500 TLD reader to measure the doses.

**Result:** The descriptive analysis showed that the mean of testis dose is directly proportional to the increase various tube potentials. The mean of ovary dose is inversely proportional to the increase various tube potentials. The mean dose of reduction percentage for ovary is 5.5% which is higher than reduction dose in testis. The mean dose of reduction percentage for testis is about 2.3%. Two-way ANOVA test for reproductive organs doses at various kVp found that there were significant difference between the variables since the p-value is  $< 0.05$  (0.01). The pos-hoc analysis (Tukey's) recommends that the p-value is  $< 0.05$  (0.01) which indicate that there is a significant difference between various tube potentials and doses in two projections.

**Conclusion:** Increase in kVp techniques significantly increases the testis dose while the ovary dose was decrease when kVp is increase. The mean dose reduction for ovary and testis was approximately about 5.5% and 2.3%. The dose reduced much higher in female compare than male when PA projection is applied to the phantom. Thus, the researcher recommends for the implementation of PA projection for routine lumbar spine examinations.

**key word:** lumbar spine, testis dose, ovary dose, AP projection, PA projection, computed radiography

# TABLE OF CONTENTS

|  |     |
|--|-----|
| AUTHOR’S DECLARATION   | ii  |
| ACKNOWLEDGEMENT  | iii |
| ABSTRACT   | iiv |
| LIST OF FIGURE   | v   |
| LIST OF TABLES   | iix |
| ABBREVIATIONS  | x   |
| <br>   |     |
| CHAPTER ONE: INTRODUCTION  | 1   |
| 1.1 Background   | 1   |
| 1.2 Problem Statement  | 2   |
| 1.3 Rationale Of The Study   | 3   |
| 1.4 Study Objectives   | 4   |
| 1.4.1 General Objective  | 4   |
| 1.4.2 Specific Objectives  | 4   |
| 1.5 Research Hypotheses  | 4   |
| <br>   |     |
| CHAPTER TWO: LITERATURE REVIEW   | 5   |
| 2.1 Introduction   | 5   |
| 2.2 Reproductive Organs Doses On Ap And Pa Positioning During Lumbar Spine Imaging | 5   |
| 2.3 Imaging Modality By Using Computed Radiography                                 | 8   |
| <br>   |     |
| CHAPTER THREE: METHODOLOGY   | 9   |
| 3.1 Introduction   | 9   |
| 3.2 Research Design  | 9   |
| 3.3 Research Setting   | 9   |
| 3.4 Scope Of Research  | 10  |
| 3.4.1 Inclusion Criteria   | 10  |

|   |        |
|---|--------|
| 3.4.2 Exclusion Criteria  | 10     |
| 3.5 Sample Selection Criteria   | 10     |
| 3.6 Data Collection Process   | 11     |
| 3.6.1 Equipments  | 11     |
| 3.6.2 Imaging of AP and PA Lumbar Spine   | 13     |
| 3.6.3 The reproductive organs doses measurement                                     | 13     |
| 3.6.4 The exposure factor for AP and PA projections                                 | 16     |
| 3.7 Statistical Consideration   | 16     |
| 3.8 Ethical Consideration   | 17     |
| <br>CHAPTER FOUR: RESULT  | <br>18 |
| 4.1 Introduction  | 18     |
| 4.2 Descriptive Analysis  | 19     |
| 4.2.2 Mean Value of Ovary Dose for Each Kvp   | 20     |
| 4.2.3 Percentage Dose Reduction for Testis and Ovary                                | 22     |
| 4.3 Two Way Anova For Each Tube Potentials  | 23     |
| 4.3.1 Two-way ANOVA test for Testis Dose in AP and PA Projection at Various kVp.    | 24     |
| <br>CHAPTER FIVE: DISCUSSION  | <br>26 |
| 5.1 Introduction  | 26     |
| 5.2 Evaluation Of Descriptive Statistic Data  | 27     |
| 5.2.1 Evaluation of Testis and Ovary dose for AP and PA Projections of Lumbar Spine | 27     |
| 5.3 Evaluation Of Two-Way Anova Test For Reproductive Organs Doses                  | 28     |
| 5.3.1 Two-way Anova Test for Testis dose in AP and PA Projections at Various kVp    | 28     |
| 5.3.2 Two-way Anova Test for Ovary dose in AP and PA Projections at Various kVp.    | 29     |
| 5.4 Justification Of Using Phantom On Present Study                                 | 29     |
| 5.5 Justification Of Using Computed Radiography                                     | 30     |
| 5.6 Weakness And Limitation   | 31     |
| <br>CHAPTER SIX: RECOMMENDATION AND CONCLUSION                                      | <br>32 |
| 6.1 Introduction  | 32     |
| 6.2 Recommendation  | 32     |
| 6.3 Conclusion  | 33     |